

## RESEARCH ARTICLE

# Bacterial and fungal symbionts of parasitic *Dendroctonus* bark beetles

Loïc Dohet<sup>1,\*</sup>, Jean-Claude Grégoire<sup>1</sup>, Aileen Berasategui<sup>2,3</sup>,  
Martin Kaltenpoth<sup>3,4</sup> and Peter H. W. Biedermann<sup>2,3</sup>

<sup>1</sup>Biological Control and Spatial Ecology Laboratory (LUBIES), CP 160/12, Université Libre de Bruxelles, av. FD Roosevelt 50, 1050 Brussels, Belgium, <sup>2</sup>Biochemistry Department, Max Planck Institute for Chemical Ecology, Hans-Knoell-Str. 8, 07745 Jena, Germany, <sup>3</sup>Insect Symbiosis Research Group, Max Planck Institute for Chemical Ecology, Hans-Knoell-Str. 8, 07745 Jena, Germany and <sup>4</sup>Department of Evolutionary Ecology, Johannes Gutenberg University of Mainz, Johann-Joachim-Becher-Weg 13, 55128 Mainz, Germany

\*Corresponding author: Biological Control and Spatial Ecology Laboratory (LUBIES), CP 160/12, Université Libre de Bruxelles, av. FD Roosevelt 50, 1050 Brussels, Belgium. Tel: +32-2-650-32-18; E-mail: [loicdohet@gmail.com](mailto:loicdohet@gmail.com)

**One sentence summary:** The parasitic bark beetles *Dendroctonus micans*, *D. punctatus* and *D. valens* house bacterial and fungal communities commonly associated with tree-inhabiting insects, and likely benefit from these symbionts in various ways.

Editor: Rolf Kümmerli

<sup>†</sup>Loïc Dohet, <http://orcid.org/0000-0002-0342-8980>

## ABSTRACT

Bark beetles (Curculionidae: Scolytinae) are one of the most species-rich herbivorous insect groups with many shifts in ecology and host-plant use, which may be mediated by their bacterial and fungal symbionts. While symbionts are well studied in economically important, tree-killing species, little is known about parasitic species whose broods develop in living trees. Here, using culture-dependent and independent methods, we provide a comprehensive overview of the associated bacteria, yeasts and filamentous fungi of the parasitic *Dendroctonus micans*, *D. punctatus* and *D. valens*, and compare them to those of other tree-inhabiting insects. Despite inhabiting different geographical regions and/or host trees, the three species showed similar microbial communities. Enterobacteria were the most prevalent bacteria, in particular *Rahnella*, *Pantoea* and *Ewingella*, in addition to *Streptomyces*. Likewise, the yeasts *Candida*/*Cyberlindnera* were the most prominent fungi. All these microorganisms are widespread among tree-inhabiting insects with various ecologies, but their high prevalence overall might indicate a beneficial role such as detoxification of tree defenses, diet supplementation or protection against pathogens. As such, our results enable comparisons of symbiont communities of parasitic bark beetles with those of other beetles, and will contribute to our understanding of how microbial symbioses facilitate dietary shifts in insects.

**Keywords:** insect symbiosis; bacteria; fungi; *Dendroctonus*; parasitic bark beetles; ecology

## INTRODUCTION

Herbivory by insects is evolutionary derived (Labandeira and Sepkoski 1993) and highly successful given the accelerated rates of speciation after switching to herbivorous diet (Mitter, Farrell

and Wiegmann 1988; Farrell 1998). Microbial symbionts play a major role in this transition, because many plant materials are nutritionally imbalanced for herbivorous insects (Watanabe and Tokuda 2010). Symbiotic bacteria and fungi can provide their

insect hosts with essential capabilities for synthesizing nutrients, overcoming plant defenses, digesting lignocellulosic plant tissue and detoxifying plant defensive chemicals (Douglas 2009; Gibson and Hunter 2010). In addition, microbial symbionts may also help herbivorous insects to protect themselves against microbial competitors, pathogens and/or predators (Flórez et al. 2015).

Bark beetles (Coleoptera: Curculionidae: Scolytinae) are one of the most successful herbivorous insect groups with at least 6000 known species worldwide (Knížek and Beaver 2004). Most of these attack woody plant tissues and studied species harbor diverse bacterial and fungal symbionts. As for other insects, these symbionts offer different services to their hosts, which researchers are just beginning to understand (Six 2013; Hofstetter et al. 2015). The tremendous diversity of bark beetle feeding habits and ecologies as well as the repeated evolutionary switches between these make bark beetle–microbe symbioses a relevant model for studying the function and evolution of insect symbionts in relation to host ecology (Six and Klepzig 2004; Six 2012; Kirkendall, Biedermann and Jordal 2015).

The relatively small genus *Dendroctonus* (19 described species) is the best studied bark beetle group today, and fungal as well as bacterial symbionts have been studied in several species (Six and Klepzig 2004; Davis 2014; Hofstetter et al. 2015). They are fascinating from an evolutionary point of view, because of their diversity of host-use strategies which are expected to strongly shape symbiont communities (Six and Klepzig 2004; Six 2012). *Dendroctonus* species have been classified in three groups depending on their host use: (i) aggressive tree killers (henceforth termed *aggressive beetles*; e.g. *Dendroctonus frontalis*, *D. ponderosae*), (ii) parasites on living trees that do not kill their host (henceforth termed *parasitic beetles*; e.g. *D. micans*, *D. punctatus*) and (iii) early successional saprophages on dead or dying trees (henceforth termed *saprophagous beetles*; e.g. *D. approximatus*, although it has also been reported to attack healthy trees) (Lindgren and Raffa 2013; Six and Bracewell 2015). As already observed by Lindgren and Raffa (2013), *D. valens* is somehow unusual, with a flexible status. In its native range, it would usually rather qualify as an early saprophage on the stumps of freshly cut hosts, or on lightning-struck trees, but it can also be parasitic on living trees (Smith 1971). Quite contrastingly, the North American *D. valens* aggressively attacks and kills living trees in China, where it was introduced and became epidemic under climatic conditions stressful for the host trees (Sun et al. 2013). On the whole, this species should thus rather be seen as *near parasitic*.

Although all *Dendroctonus* spp. invariably attack conifer hosts, these aggressive, parasitic and saprophagous species are faced with completely different challenges according to the types of hosts they colonize, and are therefore expected to profit from symbionts in different ways (Six and Klepzig 2004; Lindgren and Raffa 2013). Aggressive beetles, for example, would profit from symbionts that help them to overcome the tree defenses and to deal with declining phloem quality (Six and Klepzig 2004; Bleiker and Six 2007). By contrast, parasitic beetles are not expected to associate with symbionts that seriously harm the host tree, but instead with ones that help them to detoxify constitutive and continuously induced chemical tree defenses. The fact that only symbionts of aggressive beetles are well studied, however, currently limits our ability to develop a theoretical framework describing how host-use impacts symbiont communities and their functional roles in *Dendroctonus* (Six and Bracewell 2015).

The filamentous fungi are the best studied symbionts of *Dendroctonus*. Many aggressive beetles are obligately dependent on

fungal symbionts, which they transmit from their natal brood system to the new host in highly selective spore-carrying organs, called mycangia, on the exoskeleton (Six 2003). These mycangial fungi supplement the beetles' phloem diet with additional nitrogen and sterols (Ayres et al. 2000), and in some cases may help beetles to overcome tree defenses (Six and Klepzig 2004); although the latter role has been questioned recently (Six and Wingfield 2011). Many filamentous fungi can be isolated from bark beetles, but only those found with a high prevalence (notably in mycangia) can be regarded as obligate mutualists. The limited information available on parasitic bark beetles suggests that they do not have mycangia and do not engage in obligate mutualisms with filamentous fungi (Six and Bracewell 2015).

Yeasts are ubiquitous associates of bark beetles, but their functional role for the host is not well understood. In *Dendroctonus*, they are often isolated at high rates, but only few are host-specific (Rivera et al. 2009). They can play a beneficial role in beetle pheromone communication (Zhao et al. 2015) and some enhance the growth of fungal mutualists while suppressing fungal competitors and/or entomopathogens *in vitro* (Adams et al. 2008; Davis et al. 2013). Others negatively affect beetles by attracting natural enemies or by producing toxic chemicals (Boone et al. 2008). Again, in *Dendroctonus*, yeast communities are mostly described for the aggressive beetles.

Bacterial symbionts of *Dendroctonus* are ubiquitous in beetle guts, mycangia and galleries. Many taxa could provide nutritional benefits to the beetles by accessing sugars from complex polymers, recycling nitrogen from beetle excretions or fixing atmospheric nitrogen (Engel and Moran 2013; Morales-Jiménez et al. 2013). Other bacterial symbionts can detoxify host tree defensive compounds (Adams et al. 2013; Boone et al. 2013), defend their hosts against microbial competitors or pathogens (Cardoza, Klepzig and Raffa 2006; Scott et al. 2008) and/or facilitate the growth of the fungal mutualists (Adams et al. 2008). Generally, *Dendroctonus* species host relatively species-poor bacterial communities compared to other insects; this is likely due to the protected nature of the habitat and the toxicity of the phloem (Franceschi et al. 2005; Six and Bracewell 2015). Many symbionts of *Dendroctonus* are also known as symbionts of plants, likely because of their pre-adaptations to plant defenses. Whether indeed plant symbionts play a role in the beetles' success and what proportion of the insects' symbionts is picked up anew from the tree every generation is currently unknown (despite recent attempts, e.g. Mason, Hanshew and Raffa 2016).

*Dendroctonus micans* (Kugelann) and *D. punctatus* LeConte are two parasitic species with similar ecologies; *D. valens* LeConte seems to have a different ecology (see above). However, all three can complete their entire life cycle within living hosts: *D. micans* mainly on spruce in Europe and Asia, *D. punctatus* on spruce in Northern USA, Canada and Alaska, and *D. valens* on pines in Mexico, USA and Southern Canada (Smith 1971; Wood 1982; Grégoire 1988; Furniss 1995). A single female (*D. micans* and *D. punctatus*) or pair (*D. valens*) bores a gallery in the inner bark, often near ground level, and lays batches of eggs. Larvae feed gregariously in a communal chamber where frass is accumulated and pupation occurs. Emerging adults either mate with siblings or, in the case of *D. valens*, after dispersal. Microbial symbionts have been studied in several populations of *D. valens* (Six and Klepzig 2004; Rivera et al. 2009; Adams et al. 2010). Surveys of *D. micans*'s symbionts were limited to culturable bacteria in Turkish populations and ophiostomatoid fungi in one French population (Lieutier et al. 1992; Yilmaz et al. 2006). Non-culturable bacteria and other fungi have not been investigated in *D. micans*, and

the microbial communities of *D. punctatus* remain completely unknown.

Here we describe an extensive survey of bacterial and fungal symbionts of *D. valens*, *D. micans* and *D. punctatus* using a common set of culture-dependent and independent techniques (454 pyrosequencing). Our first objective is to characterize symbiont communities of three parasitic or near-parasitic *Dendroctonus* species. We further investigate whether these communities change during insect development or lab rearing, and if microorganisms could be taken up from the surrounding phloem. Second, we discuss how symbiont functional differences may be related to respective insect ecologies by comparing our microbial communities of parasitic bark beetles with published data on the communities of aggressive beetles.

## MATERIALS AND METHODS

### Beetle collection

#### Field insects

Mature adults and second and third instar larvae (L2-3) were sampled. *Dendroctonus micans* were collected on living trees in Commana, Merdrignac and Scrinac, Brittany, France (48°24'09.60"N 3°56'20.02"W; 48°27'51.71"N 3°38'14.97"W), in April–October 2012 and August 2013. Phloem samples were jointly taken 1 cm left or right from the edge of *D. micans* larval chambers in order to document the bacterial symbionts in the beetles' environment. *Dendroctonus punctatus* were collected on living trees west of Prince George, British Columbia, Canada (53°42'44"N 122°52'26"W; 53°43'04.57"N 122°53'07.01"W; 53°51'06"N 123°12'21"W) in July 2012 and July 2013. *Dendroctonus valens* were collected on fresh stumps and lightning-struck trees east of Redding, California, USA (40°30'28.80"N 121°51'52.77"W; 40°32'43"N 121°46'50"W; 40°43'03.78"N 121°59'27.04"W) in July 2012 and July 2013.

#### Laboratory insects

Only mature adults were sampled from the laboratory reared populations. *Dendroctonus punctatus* specimens were collected on a living tree north of Kamloops, British Columbia, Canada (51°04'29"N 120°19'52"W) in June 2011, and were reared until the second generation in the LUBIES quarantine room. *Dendroctonus valens* specimens were collected on fresh stumps in the province of Shanxi, China in 2007, and were reared until the 17th generation. As *D. micans* undergoes a reproductive diapause, laboratory adults could not be obtained for this species.

### Culture-independent bacterial community analysis

#### DNA extraction, amplification and sequencing

Samples were stored in 70% ethanol at  $-80^{\circ}\text{C}$  until analysis. Insect and phloem samples were taken out of the ethanol and dried under a sterile hood. Individual samples (whole insects or phloem) were flash frozen with liquid nitrogen and homogenized with a pestle. DNA extraction was conducted using the Epicentre MasterPure DNA Purification kit (Madison, Wisconsin, USA) in accordance with the manufacturer's instructions, including a treatment with lysozyme to break up Gram-positive bacterial cell walls. After DNA extraction, identical volumes of six individuals (except  $n = 4$  for *D. micans* field larvae and  $n = 5$  for *D. punctatus* field adults) or four phloem pieces (each  $8 \times 2 \times 2$  mm) were pooled per group for bacterial tag-encoded FLX amplicon pyrosequencing (bTEFAP). Pooled DNA samples were sent to 'Molecular Research Laboratory' (Shallowater, TX, USA) for bTEFAP with 16S rRNA primers

Gray28F forward 5'-GAGTTTGATCNTGGCTCA-3' and Gray519R reverse 5'-GTNTTACNGCGGCKGCTG-3' (Sun, Wolcott and Dowd 2011). A sequencing library was generated through one-step PCR with 30 cycles, using a mixture of HotStar and HotStar HiFidelity Taq polymerases (Qiagen, Hilden, Germany). Sequencing extended from Gray28F, using a Roche 454 FLX instrument with Titanium reagents and procedures at Molecular Research Laboratory (<http://mrdnalab.com/>).

#### 454 sequencing data analysis

QIIME (i386 1.3.0-3) was used for analysis of the 454 sequencing data (Caporaso et al. 2010). All low-quality reads (minimum average quality cut-off = 25) were removed, as well as sequences shorter than 200 bp and longer than 600 bp. No errors in the barcode but one mismatch and one ambiguous base per sequence were allowed. USEARCH6.1 (QIIME) was used to identify and discard potential chimeras. The remaining reads were analyzed using a multistep operational taxonomic unit (OTU)-picking strategy with the cdhit (Li and Godzik 2006) algorithm set at 97% similarity cut-off. The representative sequences per OTU were determined by picking the most abundant sequence, and were aligned to the Greengenes core set (<http://greengenes.lbl.gov/>) with PyNast using a minimum sequence identity of 75%. Taxonomy was assigned using the uclust classifier and only sequences with more than 0.8 confidence were selected for further analyses. Finally, an OTU table was generated with the absolute and relative abundances of bacterial phylotypes within the samples, which is visualized in a heat map constructed with MultiExperiment Viewer 4.9.0 (Saeed et al. 2003). Chloroplast and mitochondrial sequences were manually removed from the OTU table.

Alpha- and beta-diversity indices were calculated in QIIME based on the OTU table of absolute and relative abundances per sample. Rarefaction curves were generated by subsampling the OTU table with step increments of 10 sequences and 100 iterations to check for adequate sampling depth. Weighted and unweighted UniFrac distances were subjected to principal coordinates analyses (PCoA) to assess clustering of bacterial communities according to insect species, instar, phloem and laboratory-reared versus field samples. ANOSIM discriminant analyses between these groups were performed in QIIME using the unweighted and weighted UniFrac principal coordinates.

#### Phylogenetic placement of *Dendroctonus* microbiota

OTUs from the three sampled *Dendroctonus* spp. were compared with the bacterial communities of other bark beetle studies that used culture-independent methods. Our data set on *Dendroctonus*-associated bacteria was expanded by bacterial symbiont sequences of tree-inhabiting insect host species belonging to four genera (308 sequences from GenBank; Table S1, Supporting Information; see also Berasategui et al. 2016). All sequences were aligned using SINA 1.2.11 (Pruesse, Peplies and Glockner 2012), imported into ARB 5.5 (Ludwig et al. 2004) and then mapped onto the 16S SILVA rRNA database (which currently includes 597 607 curated bacterial sequences). Two to three OTUs, neighboring our OTUs, were then picked up from the SILVA database to construct a tree with our sequences and the quality checked bacterial sequences from SILVA (total 551 sequences; Table S1, Supporting Information). The quality of the alignment for each OTU sequence was manually checked and corrected. A total of 32 of the initial 182 OTUs that were imported in ARB did not fit among the 16S SILVA sequences and were excluded from our tree. The final tree was constructed with the remaining 150 OTUs in FastTree 2.1 using the GTR model (Price et al. 2010) and was edited in MEGA 6 (Tamura et al. 2011).



## Culture-dependent microbiota analysis

### Bacterial and fungal isolation

Live insects ( $n = 6$ ) were individually surface-sterilized in 70% ethanol, rinsed in distilled water and crushed as a whole in 500  $\mu\text{L}$  phosphate buffer solution with a Retsch MM301 grinder and beads (Haan, Germany). The supernatant was serially diluted 10-fold in 10 mM  $\text{MgSO}_4$  (up to  $10^{-8}$ ). Dilutions were plated ( $n = 2$ ) on potato dextrose agar (PDA), yeast extract malt agar (YEMA) and nitrogen-depleted medium described in Emtiazi, Pooyan and Shamalnasab (2007) to detect potential nitrogen-fixing bacteria. Plates were incubated at 30°C for optimal growth (1 to 14 days), and colony forming units (CFU) were counted based on plated dilutions. Pure cultures of all morphotypes were obtained and preserved in glycerol at -80°C. Yeasts were discriminated from bacteria microscopically.

### DNA extraction, amplification and sequencing

Typical 2–3 mm diameter colonies of bacteria and yeasts were transferred to a cell lysis solution (67 mM Tris-HCl pH 8.8, 16.6 mM  $(\text{NH}_4)_2\text{SO}_4$ , 5 mM  $\beta$ -mercaptoethanol, 6.7 mM  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ , 6.7  $\mu\text{M}$  EDTA pH 8.0, 1.7  $\mu\text{M}$  SDS), while filamentous fungi were grown on cellophane and ground in liquid nitrogen for DNA extraction using the MasterPure DNA Purification kit (Epicentre).

### Bacterial 16S amplification

For bacterial isolates, the 16S rRNA gene was amplified with primers fd1 (5'-AGAGTTTGTATCCTGGCTCAG-3') and rP2 (5'-ACGGCTACCTTGTTACGACTT-3') (Weisburg et al. 1991) using the following PCR conditions: 94°C for 3 min; 35 cycles of 94°C for 40 s, 60°C for 1 min, 72°C for 1 min and a final extension of 72°C for 4 min. PCR was conducted with 25 U  $\text{mL}^{-1}$  TopTaq DNA polymerase and 1x TopTaq PCR buffer (Qiagen), 0.2 mM of each dNTP, 0.2  $\mu\text{M}$  of each primer and typically 2.3–5 ng  $\mu\text{L}^{-1}$  of template DNA. PCR products were purified using 1.54 U  $\mu\text{L}^{-1}$  Exonuclease I (Epicentre) and 0.15 U  $\mu\text{L}^{-1}$  FastAP Thermosensitive Alkaline Phosphatase (ThermoFisher Scientific, Erembodegem, Belgium), and sequenced with the BigDye Terminator v3.1 Cycle Sequencing Kit on a 3730 DNA Analyzer (Applied Biosystems, Foster City, California, USA) with the same primers and additionally the primer R1087 (5'-CTCGTTGCGGCACCTTAACCC-3') for the 16S rDNA.

### Fungal 18S and ITS amplification

For fungal isolates, two genetic markers were sequenced. First, the 18S rRNA gene was amplified with the primers NS1 (5'-GTAGTCATATGCTTGCTC-3') and NS4 (5'-CTTCCGTCATATYCTTTAAG-3') (White et al. 1990) using the following PCR conditions: 94°C for 30 s; 35 cycles of 94°C for 30 s, 55°C for 45 s, 72°C for 1 min; and 72°C for 7 min. Second, the ITS region was amplified with the primers ITS1 (5'-TCCGTAGGTGAACCTGCGG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') (White et al. 1990) with PCR conditions as follows: 92°C for 2 min; 41 cycles of 95°C for 30 s, 55°C for 30 s, 72°C for 1 min; 72°C for 8 min (Six et al. 2009). PCR and sequencing mixtures were the same as for bacteria, except that primer couples NS1–NS4 and ITS1–ITS4 were used.

### BLAST

Consensus sequences were assembled in Geneious Pro 4.8.5 (Biomatters, Auckland, New Zealand) and blasted against the GenBank database. Taxa were considered identified at the species level with a match level of at least 97% identity, at the genus level with a match of at least 95%, and were considered

unidentified below 95%. Phylogenetic relationships among isolated strains and closest BLASTn matches were constructed using MOLE-BLAST with the neighbor-joining method.

## RESULTS

### Culture-independent bacterial community analysis

Our metagenomic analysis of *Dendroctonus micans*, *D. punctatus* and *D. valens* field adults, larvae, phloem and laboratory-reared adults yielded a total of 169 107 sequences (after quality trimming), which were clustered into 182 OTUs. OTUs were assigned to the level of 56 different genera, 57 families, 32 orders and 20 classes of bacteria. In the relative abundance heat map of the 50 most abundant OTUs (Fig. 1), OTUs 230 (*Luteibacter*), 124 (*Pseudomonas*), 251 (*Ralstonia*), 216 (*Erwinia*), 50 (other unidentified Xanthomonadaceae), 166 and in particular 52 (both other unidentified Enterobacteriaceae) were the most conserved among the three studied parasitic or near-parasitic bark beetles. Within *D. punctatus*, the prominent *Erwinia* (OTU 216) was almost absent from the second generation of lab-reared adults while an unidentified enterobacterium (OTU 52, close to *Rahnella* spp.) was 3.5 times as abundant as in field adults. Similarly, the rare enterobacterial OTU 123 (also close to *Rahnella* spp.) was more than 10 times as abundant and many less abundant OTUs were completely absent in the 17th generation lab-reared *D. valens*, in comparison with the field population.

Except *D. micans* field adults, whose community was predominated by Firmicutes (*Lactococcus*), all other samples mainly harbored Gammaproteobacteria (Fig. 2). Depending on the sample, these were mainly *Acinetobacter* (only in *D. micans* field larvae and nearby phloem) or *Erwinia* and other unidentified genera in all others. Betaproteobacteria (*Ralstonia* and other Comamonadaceae) were common only in *D. punctatus* and *D. micans* field larvae and nearby phloem. Rarefaction analyses demonstrated sufficient sampling depth for all our samples (Fig. S1, Supporting Information). Field-collected *D. micans* hosted less bacterial OTUs ( $25 \pm 5$  OTUs, mean  $\pm$  SD) than *D. punctatus* ( $57 \pm 23$ ) and *D. valens* ( $46 \pm 6$ ) (Table 1). Field-collected adults and larvae were more species rich ( $45 \pm 3$ ) than those reared in the laboratory ( $25 \pm 9$ ).

The weighted PCoA components explained a total of 82.8% of the variance in the OTUs' relative abundance (Fig. S2, Supporting Information). All instars of *D. punctatus* and *D. valens* clustered together, except *D. punctatus* field larvae, which were a bit closer to another cluster composed of *D. micans* field larvae and nearby phloem. *Dendroctonus micans* field adults were distant from all other clusters. The unweighted PCoA components explained 45.7% of variance (Fig. S3, Supporting Information). There was no effect of species (ANOSIM  $P = 0.71$ ) and origin ( $P = 0.42$ ) on the bacterial communities based on unweighted UniFrac. However, a significant effect of stages ( $P = 0.023$ ) on bacterial communities was found. This analysis needs to be interpreted with care since (i) abundance of OTUs lead to important differences between the weighted and unweighted PCoA plots (Figs S2 and S3, Supporting Information), and (ii) ANOSIM on weighted UniFrac values only detected a tendency for clustering for species ( $P = 0.056$ ) but not for origin ( $P = 0.39$ ) and stage ( $P = 0.39$ ).

The phylogenetic position of the three sampled *Dendroctonus* species' OTUs is shown in comparison with the bacterial community of other bark- and wood-inhabiting insects, and bacteria isolated from the environment (Table S1; Figs. 3 and S4, S5, Supporting Information). The *Dendroctonus* OTUs were close




Figure 1. Relative abundance heat map (log-transformed, white = absence, dark blue = high) of the 50 most abundant pyrosequenced bacterial OTUs (based on the cumulative relative abundance among all groups) in *D. micans*, *D. punctatus* and *D. valens* field and laboratory larvae, adults and distant phloem. Bold OTUs are the most conserved among all samples of the three beetle species, lines denote OTUs with corresponding isolates (i.e. identity > 99.3%) and accession numbers (AN) that we gained by culturing on PDA, YEMA and/or nitrogen-depleted media.




Figure 2. Relative abundance of pyrosequenced bacterial genera in *D. micans* (M), *D. punctatus* (P) and *D. valens* (V) field (w) and laboratory (l) larvae (L), adults (A) and distant phloem (Ph).



**Table 2.** Identity and prevalence (% of sampled individuals) of bacterial and fungal isolates that we gained by culturing on PDA, YEMA and nitrogen-depleted media from *D. micans*, *D. punctatus* and *D. valens* field larvae and adults. Asterisk indicates isolates that were able to grow on nitrogen-depleted medium, = = denotes synonyms, / indicates assignments equally closely related.

Prevalence (%)	<i>Dendroctonus micans</i>		<i>Dendroctonus punctatus</i>		<i>Dendroctonus valens</i>	
	Larvae	Adults	Larvae	Adults	Larvae	Adults
<b>Bacteria</b>						
<i>Enterobacter</i> sp.	0	0	0	0	0	33
<i>Erwinia billingiae</i>	0	0	0	0	0	17
<i>Erwinia</i> sp.	0	0	0	0	17	0
<i>Erwinia typographi</i>	0	0	0	33*	33*	0
<i>Ewingella</i> sp.	0	0	0	83*	33	0
<i>Pantoea agglomerans</i>	0	17*	0	0	0	67*
<i>Pantoea cedenensis</i>	0	0	67*	33*	50*	50*
<i>Pseudomonas</i> sp. 1	0	0	0	0	17*	33
<i>Pseudomonas</i> sp. 2	0	0	0	0	17*	0
<i>Rahnella aquatilis</i>	67*	100*	17*	0	83*	67
<i>Rouxiella</i> sp.	0	67*	0	0	0	0
<i>Serratia liquefaciens</i>	0	0	0	17	0	0
<i>Serratia marcescens</i>	0	0	0	0	17*	0
<i>Serratia</i> sp. 1	0	0	0	0	0	33*
<i>Serratia</i> sp. 2	0	17*	0	0	0	0
<i>Streptomyces lienomycini</i>	0	0	67	17	0	0
<b>Fungi</b>						
<i>Acanthophysium cerussatum</i> / <i>Stereum gausapatum</i>	17*	0	0	17*	0	0
<i>Candida piceae</i>	33	0	33	17	33	67
other unidentified <i>Candida</i> spp.	83*	33*	0	17*	0	67*
<i>Candida</i> sp. 1/ <i>Candida</i> sp. 3	33*	17	0	0	0	17
<i>Candida</i> sp. 2/ <i>C. fructus</i>	50*	67*	0	17*	0	0
<i>Cyberlindnera americana</i> / <i>Candida</i> sp. 3	33	17	0	0	33	17*
<i>Hypocrea muroiana</i> / <i>Trichoderma viride</i>	0	0	83*	0	0	0
<i>Penicillium charlesii</i> / <i>P. pulvis</i>	0	0	17*	0	0	17
<i>Penicillium chrysogenum</i>	0	17*	0	0	0	0
<i>Penicillium corylophilum</i>	0	17*	0	0	0	0
<i>Penicillium decumbens</i>	17	0	0	0	0	0
<i>Penicillium purpurogenum</i> = = <i>Talaromyces purpurogenus</i>	0	0	17	0	0	0
other unidentified <i>Penicillium</i> spp.	17*	0	17	17*	0	0
<i>Penicillium</i> sp. 1/ <i>Talaromyces radicus</i>	0	0	17	0	0	0
<i>Penicillium</i> sp. 2/ <i>Talaromyces variabilis</i>	17*	50*	0	0	0	17*
<i>Pichia bispora</i> /Yeast sp. 5	0	0	67	0	0	0
Fungus sp. 8	0	17	0	0	0	0
Yeast sp. 7	0	0	0	0	0	17

### Bacterial community

The mean bacterial population was  $1.7 \times 10^7$  CFUs/adult and  $6.0 \times 10^5$  CFUs/larva (Figs S6 and S7, Supporting Information; Table 2). *Rahnella aquatilis* was the most widespread bacterium; it consistently inhabited all instars of *D. micans* and *D. valens* ( $\geq 67\%$  of prevalence). Similarly, *Pantoea cedenensis* was present in at least 50% of *D. punctatus* larvae and all instars of *D. valens*. Other very prevalent taxa were *Ewingella* sp. in 83% of *D. punctatus* adults, *Streptomyces lienomycini* in 67% of *D. punctatus* larvae, *Pa. agglomerans* in 67% of *D. valens* adults and *Rouxiella* sp. in 67% of *D. micans* adults.

### Fungal community

The mean fungal population was  $4.9 \times 10^7$  CFUs/adult and  $2.9 \times 10^4$  CFUs/larva. Overall, yeasts were the most abundant fungi (Figs 4, S8 and S9, Supporting Information; Table 2), especially *Candida*/*Cyberlindnera* spp. They were isolated from all three insect species and dominated particularly in *D. micans*

and in *D. valens* field adults ( $\geq 67\%$  of samples). These yeasts were less abundant in *D. punctatus* field larvae, where *Hypocrea muroiana*/*Trichoderma viride* (83%) and *Pichia bispora*/Yeast sp. 5 (67%) were found instead. *Penicillium*/*Talaromyces* spp. were the most abundant filamentous fungi and were only absent in *D. valens* field larvae. Despite being ubiquitous, these were inconsistent associates among the insects, except *Penicillium* sp. 2/*Talaromyces variabilis* in *D. micans* field adults (50%).

### Complementarity of culture-dependent and independent analyses

Bacterial 16S sequences from cultured strains and pyrosequenced OTUs were aligned for potential matches between both methods. Seven OTUs (Fig. 1; Table S2, Supporting Information) corresponded with six different Enterobacteriaceae species (*Erwinia billingiae*, *E. typographi*, *Ewingella* sp. 1, *Pa. agglomerans*, *Ra. aquatilis*, *Serratia* sp. 2) and two Pseudomonadaceae species






Figure 4. Relative abundance of cultured fungal taxa in *D. micans* (M), *D. punctatus* (P) and *D. valens* (V) field (w) larvae (L) and adults (A).

(*Pseudomonas* sp. 1 and 2), accounting for a total of 101 305 sequences (59.9% of sequences from all OTUs).

## DISCUSSION

In this study, we characterized the bacterial and fungal communities associated with three bark beetles, *Dendroctonus micans*, *D. punctatus* and *D. valens*, which have a similar parasitic or near-parasitic ecology. Despite inhabiting distinct geographical regions and/or different host trees, these insects shared a large proportion of their microbial symbionts: (1) Enterobacteriaceae (Gammaproteobacteria) was the dominant bacterial family with the most prevalent species *Rahnella aquatilis*, *Pantoea cedenensis*, *Ewingella* sp., *Pa. agglomerans* and *Rouxiiella* sp. (although this one is very close to *Ra. aquatilis*). Enterobacteria, especially the aforementioned genera and species, are often reported as symbionts of bark- and wood-boring insects (Morales-Jiménez et al. 2009, 2012; Hu et al. 2013; Aylward et al. 2014; Xu et al. 2015; Berasategui et al. 2016; Mason, Hanshew and Raffa 2016). (2) *Streptomyces lienomycini* was commonly isolated from *D. punctatus*. *Streptomyces* bacteria are frequent associates of North American bark beetles (Hulcr et al. 2011). (3) Yeasts were the most abundant fungi, especially *Candida/Cyberlindnera* spp. in *D. micans* and *D. valens*, whereas *D. punctatus* also commonly harbored *Pichia bispora*. Likewise, these yeasts are frequent symbionts of bark and ambrosia beetles (Rivera et al. 2009; Davis 2014; Lou, Lu and Sun 2014).

All these prevalent microorganisms are close relatives of common, free-living bacteria (Figs 3, S4 and S5, Supporting Information) and yeasts. *Rahnella aquatilis*, *Ewingella* spp., *Pa. agglomerans* as well as *Candida* spp. are regularly isolated from soil, water and also from plants as epiphytes and endophytes (Weinthal et al. 2007; Winder, Macey and Cortese 2010; Ribeiro and Cardoso 2012; Hu, Li and Chen 2015). The latter means that they likely occur in the vicinity of bark- and wood-boring beetles, which may explain why they are so frequently isolated from

these insects (Colman, Toolson and Takacs-Vesbach 2012). As a striking illustration of endophyte intake, *D. micans* larvae housed bacteria very similar to nearby phloem, with prominent *Acinetobacter* and *Comamonadaceae*. Our phloem samples from just *D. micans*, however, have low explanatory power in this respect and it remains unknown which microorganisms are present in the phloem already before bark beetles establish their galleries there (Mason, Hanshew and Raffa 2016). Nevertheless, even if their associated microbes can occur in the environment, this does not exclude possible functional roles for the performance of *D. micans*, *D. punctatus* and *D. valens*. Facultative symbionts can make important contributions to host fitness, even when they are acquired from the environment (Kikuchi, Hosokawa and Fukatsu 2007; Engel and Moran 2013).

Overall, the bacterial community associated with (near-) parasitic bark beetles does not seem very constant. Unlike *D. punctatus*, the bacterial communities of *D. valens* and *D. micans* have already been studied before and greatly vary geographically (Yilmaz et al. 2006; Morales-Jiménez et al. 2009; Adams et al. 2010; Xu et al. 2015; this study). In *D. valens*, the bacterial community from the Californian population we sampled (where *D. valens* qualifies as a saprophagous-parasitic species) was much more distant to the one from the Northern USA (Adams et al. 2010) than to the ones from Mexico (Morales-Jiménez et al. 2009) or from invasive populations in China (although *D. valens* behaves aggressively there; Xu et al. 2015). Despite some geographical variation, *Rahnella* and *Pantoea* were consistent symbionts throughout all four studies. In the bacterial community of *D. micans*, there is only a small overlap (mainly *Serratia*) in the identified taxa from France (this study) and from Turkey (Yilmaz et al. 2006). By contrast to these parasitic beetles, the bacterial community of the aggressive *D. ponderosae* is much more stable (Adams et al. 2013). Variable bacterial communities could be related to rather opportunistic, loose symbiotic associations, with little effect for their host (Engel and Moran 2013). However, symbiont redundancy, where different microorganisms play similar roles (Six 2012), could also explain this pattern. Throughout



the developmental stages, the bacterial community of *D. valens* was very constant, but the ones of both *D. micans* and *D. punctatus* were quite distinct between field-collected adults and larvae. Several non-exclusive hypotheses could explain such community changes during the developmental stages: (i) successive molts disrupt the gut bacterial flora (Engel and Moran 2013); (ii) adults and larvae feed on different parts of the phloem, which may be of different quality (e.g. in terms of nutrition and defense compounds; Franceschi et al. 2005); (iii) larvae and adults select different symbionts according to their needs; (iv) temporal turnover of the symbionts (unlike *D. punctatus*, *D. micans* larvae and adults were collected in the same place but at different times).

The bacterial species richness was much lower in laboratory-reared adults in comparison with field insects. In addition, we observed differences in the relative abundance of several bacterial OTUs. These results are similar to those of Meeus et al. (2015) who observed that *Bombus terrestris* reared indoors had a subset of the bacterial community compared to field insects. It is possible that the differences in the bacterial communities result from the rearing conditions and could, in turn, affect the beetles' physiology. In many insects, the gut community is strongly influenced by the diet (Engel and Moran 2013). In our laboratory conditions, insects were reared on fresh logs and phloem sandwiches that are likely to differ from a standing, living tree (i.e. moisture content, toxic plant compounds and general food quality; Klepzig and Six 2004; Six and Klepzig 2004) and which different microorganisms may exploit. Alternatively, in the specific case of *D. valens*, the community change between field and lab adults may be related to the different collection places.

The fungal community of *D. micans*, *D. punctatus* and *D. valens* was dominated by widespread environmental yeasts and some ubiquitous filamentous fungi. Not a single ophiostomatoid fungus was isolated. Most aggressive *Dendroctonus* spp. are tightly associated with these fungi (Six and Bracewell 2015). In its native range, *D. valens* behaves like a saprophage and parasite, and is occasionally associated with ophiostomatoid fungi, although with high variability of prevalence (Six and Klepzig 2004, Sun et al. 2013). In China, where *D. valens* aggressively attacks living trees, it is associated with phytopathogenic strains of *Lepographium procerum* and other ophiostomatoid species, however, with a very low prevalence (Lu et al. 2009, 2010; Sun et al. 2013). These insect–fungus interactions are very complex: *D. valens* could benefit from symbiotic nutritional supplementation or help in overcoming the tree defenses (Sun et al. 2013), but feeding experiments demonstrated that *L. procerum*, and other ophiostomatoid associates of *D. valens*, compete with the insects for polysaccharides in the phloem and trigger an immune response in the beetles, which resulted in lower weight gain in larvae (Shi et al. 2012; Wang et al. 2013). Such a competition for the available resources might be one reason explaining why true parasitic beetles are not consistently associated with ophiostomatoid fungi specifically, or filamentous fungi in general, which in this case are regarded as opportunistic associates (Lieutier et al. 1992; Raffa, Phillips and Salom 1993; Six and Bracewell 2015). Another reason may be the secondary metabolites of living trees that are toxic to both insects and fungi (Raffa and Smalley 1995; Franceschi et al. 2005; Krokene 2015), but to which parasitic beetles evolved a high resistance (Everaerts, Grégoire and Merlin 1988). Last, it is also counteradaptive for parasitic beetles that derive large benefits from their life in living trees (i.e. a stable environment protecting them from generalist natural enemies and from competitors) to inoculate their galleries with fungi that have been shown (i) to trigger higher induced tree defenses,

which harm the beetles (Raffa and Smalley 1995), or even (ii) kill the tree. Future surveys should focus on the prevalence and possible roles of ophiostomatoid associates of parasitic bark beetles, which could in turn improve our knowledge on those of aggressive bark beetles.

Parasitic bark beetles live in and feed on phloem that is loaded with constitutive and induced plant defensive compounds. Therefore, symbionts that are able to assist in detoxification should be very beneficial to the beetles, like *Ra. aquatilis* (Boone et al. 2013; Xu et al. 2015) and *Candida/Cyberlindnera* spp. (Rivera et al. 2009; Lou, Lu and Sun 2014) which we both isolated at high rates in our study. By-products of such microorganisms may also play a role in the intra- or interspecific chemical communication of the beetles (Boone et al. 2008; Zhao et al. 2015). Bark beetles could also benefit from symbionts by protection and improvement of their nutritional niche. Living phloem is lacking many nutrients, like nitrogen sources and sterols (Merrill and Cowling 1966; Bentz and Six 2006). In this context, associated microorganisms could provide a significant advantage to parasitic bark beetles. First, several prevalent bacteria were isolated on nitrogen-depleted medium in this study and are known as nitrogen fixing (diazotroph): *Ra. aquatilis* (Vasanthakumar et al. 2006; Morales-Jiménez et al. 2009) and *Pa. agglomerans* (Bridges 1981). Moreover, *NifH* genes were amplified from the microbial community of *D. micans*, *D. punctatus* and *D. valens* (Dohet and Biedermann, unpublished data). Likewise, *Ra. aquatilis* is also reported as nitrogen recycling (uricolytic; Morales-Jiménez et al. 2013). Second, cellulolytic microorganisms were detected in preliminary tests on Congo red agar (Dohet, unpublished data), which may enrich the beetles' phloem diet with additional free carbohydrates. Furthermore, microorganisms such as the isolated actinomycetes *Micrococcus luteus* or *Streptomyces* spp. could shape beetles' microbial communities by producing antibiotics as shown for strains of these taxa in *D. rufipennis* and *D. frontalis*, respectively (Cardoza, Klepzig and Raffa 2006; Scott et al. 2008).

Our study suggests that parasitic bark beetles lack consistent association with highly phytopathogenic fungi but could benefit, at the same time, from other symbionts. It also discloses striking differences between parasitic bark beetles, two of which (*D. micans* and *D. punctatus*) almost never kill their hosts and are inconsistently found associated with ophiostomatoid fungi (*D. micans*; Lieutier et al. 1992) or have never been studied so far in this respect (*D. punctatus*). On the other hand, some near-parasitic bark beetles, like *D. valens*, sometimes massively kill trees and harbor ophiostomatoid fungi with a low prevalence in this case (Lu et al. 2009). *Dendroctonus murrayanae*, which also qualifies as parasitic, has been observed killing trees (Wood 1982) and is consistently associated with ophiostomatoid fungi (Six et al. 2011). In summary, we believe that these important differences in the ecology of parasitic beetles call for a substantial revision of this whole group, distinguishing true parasites from near parasites.

This is the first comprehensive characterization of the bacterial and fungal symbionts of the (near-) parasitic bark beetles *D. micans*, *D. punctatus* and *D. valens*, using a combination of culture-dependent and independent methods. Many of the close relatives of the identified taxa have been previously characterized as free-living microorganisms, but others are known as symbionts of plants, bark beetles and other wood-boring insects. Parasitic and near-parasitic bark beetles could benefit from these symbionts in various ways, through detoxification of tree defenses, diet supplementation and/or protection against pathogens. Due to the high variability of associated bacteria and fungi, future studies should sample

all developmental stages of parasitic bark beetles more intensively from various populations and at different times to identify a potential core microbiota, spatiotemporal patterns and their concrete effects on their hosts. This sampling should also take the surrounding phloem and wood into account, which will help to clarify which microbes are present within the tree beforehand and are not inoculated by the beetles. Such studies will help to understand the roles of symbionts in shifts of ecologies and hosts in bark beetles, which are a great model for the development of a broad theoretical framework on the function and evolution of bacterial and fungal symbionts in insects as a whole.

## SUPPLEMENTARY DATA

Supplementary data are available at FEMSEC online.

## ACKNOWLEDGEMENTS

The authors are grateful to Prof. Staffan Lindgren, Luke Spooner, Rylee Isitt, Chelsea Monell, Jordan Lindgren, Prof. Allan Carroll, Debra Wytrykush, Dr Nancy Gillette, Dr Tom Smith, Dr Donald Owen and Antoine Boullis for their help in the field. The authors also thank Séverine Hasbroucq and Jean-Marc Molenberg for their technical assistance, Prof. Corinne Vander Wauven for the lab training and the identification of doubtful strains, Prof. Isabelle George for advising, and Dr Celia Boone for her constructive comments on the manuscript. LD and PHWB share the corresponding authorship of this manuscript.

## FUNDING

This work was supported by the Fonds pour la Formation à la Recherche dans l'Industrie et l'Agriculture (FRIA) and travel grants were provided by the Belgian Fund for Scientific Research (FRS-FNRS). LD was supported by a doctoral grant from the FRIA and by an award from the Fonds David and Alice Van Buuren. PHWB was supported by the Swiss National Science Foundation [P300P3.151134] and by a Marie Curie Intra-European Fellowship (IEF) [project number 626279]. AB and MK acknowledge funding from the Max Planck Society.

**Conflict of interest.** None declared.

## REFERENCES

- Adams AS, Adams SM, Currie CR *et al.* Geographic variation in bacterial communities associated with the red turpentine beetle (Coleoptera: Curculionidae). *Environ Entomol* 2010;**39**:406–14.
- Adams AS, Aylward FO, Adams SM *et al.* Mountain pine beetles colonizing historical and naive host trees are associated with a bacterial community highly enriched in genes contributing to terpene metabolism. *Appl Environ Microb* 2013;**79**:3468–75.
- Adams AS, Six DL, Adams SM *et al.* *In vitro* interactions between yeasts and bacteria and the fungal symbionts of the mountain pine beetle (*Dendroctonus ponderosae*). *Microb Ecol* 2008;**56**:460–66.
- Aylward FO, Suen G, Biedermann PHW *et al.* Convergent bacterial microbiotas in the fungal agricultural systems of insects. *mBio* 2014;**5**:e02077.
- Ayres MP, Wilkens RT, Ruel JJ *et al.* Nitrogen budgets of phloem-feeding bark beetles with and without symbiotic fungi. *Ecol* 2000;**81**:2198–10.
- Bentz BJ, Six DL. Ergosterol content of fungi associated with *Dendroctonus ponderosae* and *Dendroctonus rufipennis* (Coleoptera: Curculionidae, Scolytinae). *Ann Entomol Soc Am* 2006;**99**:189–94.
- Berasategui A, Axelsson K, Schmidt A *et al.* The gut microbiota of the pine weevil is similar across Europe and resembles that of other conifer-feeding beetles. *Mol Ecol* 2016, DOI:10.1111/mec.13702.
- Bleiker KP, Six DL. Dietary benefits of fungal associates to an eruptive herbivore: potential implications of multiple associates on host population dynamics. *Environ Entomol* 2007;**36**:1384–96.
- Boone CK, Keefover-Ring K, Mapes AC *et al.* Bacteria associated with a tree-killing insect reduce concentrations of plant defense compounds. *J Chem Ecol* 2013;**39**:1003–06.
- Boone CK, Six DL, Zheng Y *et al.* Parasitoids and dipteran predators exploit volatiles from microbial symbionts to locate bark beetles. *Environ Entomol* 2008;**37**:150–61.
- Bridges J. Nitrogen-fixing bacteria associated with bark beetles. *Microb Ecol* 1981;**7**:131–7.
- Caporaso JG, Kuczynski J, Stombaugh J *et al.* QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* 2010;**7**:335–6.
- Cardoza YJ, Klepzig KD, Raffa KF. Bacteria in oral secretions of an endophytic insect inhibit antagonistic fungi. *Ecol Entomol* 2006;**31**:636–45.
- Colman DR, Toolson EC, Takacs-Vesbach CD. Do diet and taxonomy influence insect gut bacterial communities? *Mol Ecol* 2012;**21**:5124–37.
- Davis TS. The ecology of yeasts in the bark beetle holobiont: a century of research revisited. *Microb Ecol* 2014;**69**:723–32.
- Davis TS, Crippen TL, Hofstetter RW *et al.* Microbial volatile emissions as insect semiochemicals. *J Chem Ecol* 2013;**39**:840–59.
- Douglas A. The microbial dimension in insect nutritional ecology. *Funct Ecol* 2009;**23**:38–47.
- Emtiazi G, Pooyan M, Shamalnasab M. Cellulase activities in nitrogen fixing *Paenibacillus* isolated from soil in N-free media. *World J Agr Sci* 2007;**3**:602–8.
- Engel P, Moran NA. The gut microbiota of insects - diversity in structure and function. *FEMS Microbiol Rev* 2013;**37**:699–735.
- Everaerts C, Grégoire J-C, Merlin J. The toxicity of Norway spruce monoterpenes to two bark beetle species and their associates. In: Mattson WJ, Levieux J, Bernard-Dagan C (eds). *Mechanism of Woody Plant Defenses Against Insects*. New York: Springer, 1988, 335–44.
- Farrell BD. 'Inordinate fondness' explained: why are there so many beetles? *Science* 1998;**281**:555–59.
- Flórez LV, Biedermann PHW, Engl T *et al.* Defensive symbioses of animals with prokaryotic and eukaryotic microorganisms. *Nat Prod Rep* 2015;**32**:904–36.
- Franceschi VR, Krokene P, Christiansen E *et al.* Anatomical and chemical defenses of conifer bark against bark beetles and other pests. *New Phytol* 2005;**167**:353–75.
- Furniss MM. Biology of *Dendroctonus punctatus* (Coleoptera: Scolytidae). *Ann Entomol Soc Am* 1995;**88**:173–82.
- Gibson CM, Hunter MS. Extraordinarily widespread and fantastically complex: comparative biology of endosymbiotic bacterial and fungal mutualists of insects. *Ecol Lett* 2010;**13**:223–34.
- Grégoire J-C. The greater European spruce beetle. In: Berryman AA (ed). *Dynamics of Forest Insect Populations: Patterns, Causes, Implications*. New York: Springer Science+Business Media, 1988, 455–78.

- Hofstetter RW, Dinkins-Bookwalter J, Davis TS et al. Symbiotic associations of bark beetles. In: Vega FE, Hofstetter RW (eds). *Bark Beetles*. San Diego: Academic Press, 2015, 209–45.
- Hu X, Li M, Chen H. Community structure of gut fungi during different developmental stages of the Chinese white pine beetle (*Dendroctonus armandi*). *Sci Rep* 2015;5:8411.
- Hu X, Wang C, Chen H et al. Differences in the structure of the gut bacteria communities in development stages of the Chinese white pine beetle (*Dendroctonus armandi*). *Int J Mol Sci* 2013;14:21006–20.
- Hulcr J, Adams AS, Raffa K et al. Presence and diversity of *Streptomyces* in *Dendroctonus* and sympatric bark beetle galleries across North America. *Microb Ecol* 2011;61:759–68.
- Kikuchi Y, Hosokawa T, Fukatsu T. Insect-microbe mutualism without vertical transmission: a stinkbug acquires a beneficial gut symbiont from the environment every generation. *Appl Environ Microb* 2007;73:4308–16.
- Kirkendall LR, Biedermann PHW, Jordal BH. Evolution and diversity of bark and ambrosia beetles. In: Vega FE, Hofstetter RW (eds). *Bark Beetles*. San Diego: Academic Press, 2015, 85–156.
- Klepzig KD, Six DL. Bark beetle-fungal symbiosis: context dependency in complex associations. *Symbiosis* 2004;37:189–205.
- Knížek M, Beaver R. Taxonomy and systematics of bark and ambrosia beetles. In: Lieutier F, Day KR, Battisti A et al. (eds). *Bark and Wood Boring Insects in Living Trees in Europe, a Synthesis*. Dordrecht: Springer, 2004, 41–54.
- Krokene P. Conifer defense and resistance to bark beetles. In: Vega FE, Hofstetter RW (ed.). *Bark Beetles*. San Diego: Academic Press, 2015, 177–207.
- Labandeira CC, Sepkoski JJ. Insect diversity in the fossil record. *Science* 1993;261:310–15.
- Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* 2006;22:1658–9.
- Lieutier F, Vouland G, Pettinetti M et al. Defence reactions of Norway spruce (*Picea abies* Karst.) to artificial insertion of *Dendroctonus micans* Kug. (Col., Scolytidae). *J Appl Entomol* 1992;114:174–86.
- Lindgren BS, Raffa KF. Evolution of tree killing in bark beetles (Coleoptera: Curculionidae): trade-offs between the maddening crowds and a sticky situation. *Can Entomol* 2013;145:471–95.
- Lou Q-Z, Lu M, Sun J-H. Yeast diversity associated with invasive *Dendroctonus valens* killing *Pinus tabulaeformis* in China using culturing and molecular methods. *Microb Ecol* 2014;68:397–15.
- Lu M, Wingfield MJ, Gillette NE et al. Complex interactions among host pines and fungi vectored by an invasive bark beetle. *New Phytol* 2010;187:859–66.
- Lu Q, Decock C, Zhang XY et al. Ophiostomatoid fungi (Ascomycota) associated with *Pinus tabulaeformis* infested by *Dendroctonus valens* (Coleoptera) in northern China and an assessment of their pathogenicity on mature trees. *Anton Leeuw* 2009;96:275–93.
- Ludwig W, Strunk O, Westram R et al. ARB: a software environment for sequence data. *Nucleic Acids Res* 2004;32:1363–71.
- Mason CJ, Hanshew AS, Raffa KF. Contributions by host trees and insect activity to bacterial communities in *Dendroctonus valens* (Coleoptera: Curculionidae) galleries, and their high overlap with other microbial assemblages of bark beetles. *Environ Entomol* 2016;45:348–56.
- Meeus I, Parmentier L, Billiet A et al. 16S rRNA amplicon sequencing demonstrates that indoor-reared bumblebees (*Bombus terrestris*) harbor a core subset of bacteria normally associated with the wild host. *Plos One* 2015;10:e0125152.
- Merrill W, Cowling EB. Role of nitrogen in wood deterioration: amounts and distribution of nitrogen in tree stems. *Can J Bot* 1966;44:1555–80.
- Mitter C, Farrell B, Wiegmann B. The phylogenetic study of adaptive zones: has phytophagy promoted insect diversification? *Am Nat* 1988;132:107–28.
- Morales-Jiménez J, Vera-Ponce de León A, García-Domínguez A et al. Nitrogen-fixing and uricolytic bacteria associated with the gut of *Dendroctonus rhizophagus* and *Dendroctonus valens* (Curculionidae: Scolytinae). *Microb Ecol* 2013;66:200–10.
- Morales-Jiménez J, Zúñiga G, Ramírez-Saad HC et al. Gut-associated bacteria throughout the life cycle of the bark beetle *Dendroctonus rhizophagus* Thomas and Bright (Curculionidae: Scolytinae) and their cellulolytic activities. *Microb Ecol* 2012;64:268–78.
- Morales-Jiménez J, Zúñiga G, Villa-Tanaca L et al. Bacterial community and nitrogen fixation in the red turpentine beetle, *Dendroctonus valens* LeConte (Coleoptera: Curculionidae: Scolytinae). *Microb Ecol* 2009;58:879–91.
- Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately maximum-likelihood trees for large alignments. *PLoS One* 2010;5:e9490.
- Pruesse E, Peplies J, Glockner FO. SINA: Accurate high-throughput multiple sequence alignment of ribosomal RNA genes. *Bioinformatics* 2012;28:1823–9.
- Raffa KF, Phillips TW, Salom SM. Strategies and mechanisms of host colonization by bark beetles. In: Schowalter TD, Filip GM (eds). *Beetle-Pathogen Interactions in Conifer Forests*. London: Academic Press, 1993, 103–28.
- Raffa KF, Smalley EB. Interaction of pre-attack and induced monoterpene concentrations in host conifer defense against bark beetle-fungal complexes. *Oecologia* 1995;102:285–95.
- Ribeiro CM, Cardoso EJBN. Isolation, selection and characterization of root-associated growth promoting bacteria in Brazil Pine (*Araucaria angustifolia*). *Microbiol Res* 2012;167:69–78.
- Rivera FN, González E, Gómez Z et al. Gut-associated yeast in bark beetles of the genus *Dendroctonus* Erichson (Coleoptera: Curculionidae: Scolytinae). *Biol J Linn Soc* 2009;98:325–42.
- Saeed AI, Sharov V, White J et al. TM4: A free, open-source system for microarray data management and analysis. *BioTechniques* 2003;34:374–8.
- Scott JJ, Oh DC, Yuceer MC et al. Bacterial protection of beetle-fungus mutualism. *Science* 2008;322:63.
- Shi Z-H, Wang B, Clarke SR et al. Effect of associated fungi on the immunocompetence of red turpentine beetle larvae, *Dendroctonus valens* (Coleoptera: Curculionidae: Scolytinae). *Insect Sci* 2012;19:579–84.
- Six DL. Bark beetle-fungus symbioses. In: Bourtzis K, Miller TA (eds). *Insect Symbiosis*. Boca Raton: CRC Press, 2003, 97–114.
- Six DL. Ecological and evolutionary determinants of bark beetle—fungus symbioses. *Insects* 2012;3:339–66.
- Six DL. The bark beetle holobiont: why microbes matter. *J Chem Ecol* 2013;39:989–1002.
- Six DL, Bracewell R. *Dendroctonus*. In: Vega FE, Hofstetter RW (ed.). *Bark Beetles*. San Diego: Academic Press, 2015, 305–50.
- Six DL, de Beer ZW, Duong TA et al. Fungal associates of the lodgepole pine beetle, *Dendroctonus murrayanae*. *Anton Leeuw* 2011;100:231–44.
- Six DL, Doug Stone W, de Beer ZW et al. *Ambrosiella beaveri*, sp. nov., associated with an exotic ambrosia beetle, *Xylosandrus mutilatus* (Coleoptera: Curculionidae, Scolytinae), in Mississippi, USA. *Anton Leeuw Int J G* 2009;96:17–29.

- Six D, Klepzig K. *Dendroctonus* bark beetles as model systems for studies on symbiosis. *Symbiosis* 2004;**37**:207–32.
- Six DL, Wingfield MJ. The role of phytopathogenicity in bark beetle-fungus symbioses: a challenge to the classic paradigm. *Annu Rev Entomol* 2011;**56**:255–72.
- Smith RH. *Red Turpentine Beetle*. US Department of Agriculture, Forest Service, Forest Pest Leaflet 1971, 1–9.
- Sun J, Lu M, Gillette NE et al. Red turpentine beetle: innocuous native becomes invasive tree killer in China. *Annu Rev Entomol* 2013;**58**:293–311.
- Sun Y, Wolcott RD, Dowd SE. Tag-encoded FLX amplicon pyrosequencing for the elucidation of microbial and functional gene diversity in any environment. *Methods Mol Biol* 2011;**733**:129–41.
- Tamura K, Peterson D, Peterson N et al. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 2011;**28**:2731–9.
- Vasanthakumar A, Delalibera I, Handelsman J et al. Characterization of gut-associated bacteria in larvae and adults of the Southern pine beetle, *Dendroctonus frontalis* Zimmermann. *Environ Entomol* 2006;**35**:1710–7.
- Wang B, Lu M, Cheng C et al. Saccharide-mediated antagonistic effects of bark beetle fungal associates on larvae. *Biol Lett* 2013;**9**:20120787.
- Watanabe H, Tokuda G. Cellulolytic systems in insects. *Annu Rev Entomol* 2010;**55**:609–32.
- Weinthal DM, Barash I, Panijel M et al. Distribution and replication of the pathogenicity plasmid pPATH in diverse populations of the gall-forming bacterium *Pantoea agglomerans*. *Appl Environ Microb* 2007;**73**:7552–61.
- Weisburg WG, Barns SM, Pelletier DA et al. 16S ribosomal DNA amplification for phylogenetic study. *J Bacteriol* 1991;**173**:697–703.
- White TJ, Bruns T, Lee S et al. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ et al. (eds). *PCR Protocols: A Guide to Methods and Applications*. San Diego: Academic Press, 1990, 315–22.
- Winder RS, Macey DE, Cortese J. Dominant bacteria associated with broods of mountain pine beetle, *Dendroctonus ponderosae* (Coleoptera: Curculionidae, Scolytinae). *J Entomol Soc Brit Columbia* 2010;**107**:43–56.
- Wood SL. *The Bark and Ambrosia Beetles of North America (Coleoptera: Scolytidae), a Taxonomic Monograph*. Provo: Brigham Young University, 1982.
- Xu L, Lou Q, Cheng C et al. Gut-associated bacteria of *Dendroctonus valens* and their involvement in verbenone production. *Microb Ecol* 2015;**70**:1012–23.
- Yilmaz H, Sezen K, Kati H et al. The first study on the bacterial flora of the European spruce bark beetle, *Dendroctonus micans* (Coleoptera: Scolytidae). *Biologia (Bratisl)* 2006;**61**:679–86.
- Zhao T, Axelsson K, Krokene P et al. Fungal symbionts of the spruce bark beetle synthesize the beetle aggregation pheromone 2-methyl-3-buten-2-ol. *J Chem Ecol* 2015;**41**:848–52.